Section 1 Electric Charge

Objectives

- Understand the basic properties of electric charge.
- Differentiate between conductors and insulators.
- **Distinguish** between charging by contact, charging by induction, and charging by polarization.
- Calculate force exerted on charged particles



Resources


Properties of Electric Charge

- - like charges repel
 - unlike charges attract

- Positively charged particles are called _____
- Uncharged particles are called ______.
- Negatively charged particles are called _

Chapter menu

Resources



Properties of Electric Charge, continued

 Electric charge is _____. That is, when an object is charged, its charge is always a multiple of a _____.

Charge is measured in _____.

• The _____, is the magnitude of the charge of a single electron or proton.

or

Chapter menu

Copyright © by Holt, Rinehart and Winston. All rights reserved.

Resources

.....

Quantity of charge practice problem

• A charge of 50 elementary particles is equal to what coulomb charge?

– Where..

 $1 \text{ C} = 6.25 \text{ x} 10^{18}$ elementary charge

Chapter menu

Resources

Practice problem #2

An object *cannot* have a charge of A) 4.5 × 10⁻¹⁹ C B) 3.2 × 10⁻¹⁹ C C) 8.0 × 10⁻¹⁹ C D) 9.6 × 10⁻¹⁹ C

Chapter menu

Resources

Copyright © by Holt, Rinehart and Winston. All rights reserved.

.....

.........


0



Section 1 Electric Charge

•

The Milikan Experiment






Section 1 Electric Charge

........

Milikan's Oil Drop Experiment





Transfer of Electric Charge

- An ______ is a material in which charges can move freely.
- An ______ is a material in which charges cannot move freely.



Resources

Transfer of Electric Charge, *continued*

- Insulators and conductors can be charged by _
- Conductors can be charged by ____


is a process of charging a conductor by bringing it near another charged object and grounding the conductor.

Chapter menu


Resources

How does a electroscope respond to a charged object?

H



Chapter menu



Charging by Conduction

When charging something by contact it is important to note the following properties

The objects must actually touch and transfer some electrons.

The objects become charged alike.


The original charged object becomes less charged because it actually lost some charge. Therefore, there is a limit to how many times it could be used to charge something without being recharged.

Chapter menu

Resources

......

Charging by Induction



Charged object **does not touch** the electroscope.


Electroscope ends up **oppositely charged** to the object used to charge it.

The first charge is strong and **stays strong** each time the electroscope is recharged. (This is due to the original object not losing any charge in the process.)

Chapter menu

Resources

Transfer of Electric Charge, continued



• A surface charge can be induced on insulators by

With _____, the charges within individual molecules are realigned such that the molecule has a slight charge separation.

Chapter menu


Resources

Of

DEMO: Charge is transferred between objects

 Two spheres on glass insulating rods have the charges shown below. If the make contact and a separated, what is the charge on each sphere?

+4 C +6 C



Chapter menu


Resources

.....

0

SAMPLE PROBLEM

The diagram below shows the initial charges and positions of three metal spheres, R, S, and T, on insulating stands.



Sphere R is brought into contact with sphere S and then removed. Then sphere S is brought into contact with sphere T and removed. What is the charge on sphere T after this procedure is completed?

Chapter menu

Resources

Section 2 Electric Force


Coulomb's Law

 Two charges near one another exert a force on one another called the _____.

states that the electric force is proportional to the magnitude of each charge and inversely proportional to the square of the distance between them.

Chapter menu

Resources



......

Coulomb's Law practice problem

 What is the electrostatic force between two small spheres possessing net charges of +2.0 coulombs and -3.0 coulombs, if the distance between them is 10.0 m?



k = electrostatic constant = $8.99 \times 10^9 \text{ N} \cdot \text{m}^2 / \text{C}^2$

Chapter menu

Resources